

Representation Theory of Quantized Enveloping Algebras with Interpolating Real Structure*

Kenny DE COMMER

Department of Mathematics, University of Cergy-Pontoise,
UMR CNRS 8088, F-95000 Cergy-Pontoise, France

E-mail: Kenny.De-Commer@u-cergy.fr

URL: <http://kdecommer.u-cergy.fr>

Received August 18, 2013, in final form December 18, 2013; Published online December 24, 2013

<http://dx.doi.org/10.3842/SIGMA.2013.081>

Abstract. Let \mathfrak{g} be a compact simple Lie algebra. We modify the quantized enveloping $*$ -algebra associated to \mathfrak{g} by a real-valued character on the positive part of the root lattice. We study the ensuing Verma module theory, and the associated quotients of these modified quantized enveloping $*$ -algebras. Restricting to the locally finite part by means of a natural adjoint action, we obtain in particular examples of quantum homogeneous spaces in the operator algebraic setting.

Key words: compact quantum homogeneous spaces; quantized universal enveloping algebras; Hopf–Galois theory; Verma modules

2010 Mathematics Subject Classification: 17B37; 20G42; 46L65

Introduction

This paper reports on preliminary work related to the quantization of non-compact semi-simple Lie groups. The main idea behind such a quantization is based on the reflection technique developed in [5] and [11] (see also [7] and [6] for concrete, small-dimensional examples relevant to the topic of this paper). Briefly, this technique works as follows. Let \mathbb{G} be a compact quantum group acting on a compact quantum homogeneous space \mathbb{X} . Assume that the von Neumann algebra $\mathcal{L}^\infty(\mathbb{X})$ associated to \mathbb{X} is a type I factor. Then the action of \mathbb{G} on $\mathcal{L}^\infty(\mathbb{X})$ can be interpreted as a projective representation of \mathbb{G} , and one can deform \mathbb{G} with the ‘obstruction’ associated to this projective representation to form a new *locally* compact quantum group \mathbb{H} . More generally, if $\mathcal{L}^\infty(\mathbb{X})$ is only a finite direct sum of type I -factors, one can construct \mathbb{H} as a locally compact quantum groupoid (of a particularly simple type). Our idea is to fit the quantizations of non-compact semi-simple Lie groups into this framework, obtaining them as a reflection of the quantization of their compact companion. For this, one needs the proper quantum homogeneous spaces to feed the machinery with.

It is natural to expect the needed quantum homogeneous space to be a quantization of a compact symmetric space associated to the non-compact semi-simple Lie group. By now, there is much known on the quantization of symmetric spaces (see [25, 26] and references therein, and [32] for the non-compact situation), but these results are mostly of an algebraic nature, and not much seems known about corresponding operator algebraic constructions except for special cases. In fact, in light of the motivational material presented in Appendix B, we will instead of symmetric spaces use certain quantizations of (co)adjoint orbits, following the approach of [10, 19, 29]. Here, one rather constructs quantum homogeneous spaces as subquotients of (quantized)

*This paper is a contribution to the Special Issue on Noncommutative Geometry and Quantum Groups in honor of Marc A. Rieffel. The full collection is available at <http://www.emis.de/journals/SIGMA/Rieffel.html>

universal enveloping algebras in certain highest weight representations. We will build on this approach by combining it with real structures and the contraction technique.

Our main result, Theorem 3.20, will consist in showing that the compact quantum homogeneous spaces that we build do indeed consist of finite direct sums of type I factors. This will give a theoretical underpinning and motivation for the claim that the above mentioned quantizations of non-compact semi-simple groups can indeed be constructed using the reflection technique. Our results are however quite incomplete as of yet, as

- in the non-contracted case, we can only treat concretely the case of Hermitian symmetric spaces,
- a more detailed analysis of the resulting quantum homogeneous spaces is missing,
- the relation to known quantum homogeneous spaces is not elucidated,
- no precise connection with deformation quantization is provided,
- the relation with the approach of Korogodsky [24] towards the quantization of non-compact Lie groups remains to be clarified.

We hope to come back to the above points in future work.

The structure of this paper is as follows. In Section 1, we introduce the ‘modified’ quantized universal enveloping algebras we will be studying, and state their main properties in analogy with the ordinary quantized universal enveloping algebras. In Section 2, we introduce a theory of Verma modules, and study the associated unitarization problem. In Section 3, we study subquotients of our generalized quantized universal enveloping algebras, and show how they give rise to C^* -algebraic quantum homogeneous spaces whose associated von Neumann algebras are direct sums of type I factors. In Section 4, we briefly discuss a case where the associated von Neumann algebra is simply a type I-factor itself.

In the appendices, we give some further comments on the structures appearing in this paper. In Appendix A, we recall the notion of cogroupoids [2] which is very convenient for our purposes. In Appendix B, we discuss the Lie algebras which are implicitly behind the constructions in the main part of the paper.

1 Two-parameter deformations of quantized enveloping algebras

Let \mathfrak{g} be a complex simple Lie algebra of rank l , with fixed Cartan subalgebra \mathfrak{h} and Cartan decomposition $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$. Let $\Delta \subseteq \mathfrak{h}^*$ be the associated finite root system, Δ^+ the set of positive roots, and $\Phi^+ = \{\alpha_r \mid r \in I\}$ the set of simple positive roots. We identify I and Φ^+ with the set $\{1, \dots, l\}$ whenever convenient. Let $\mathfrak{h}_{\mathbb{R}}^* \subseteq \mathfrak{h}^*$ be the real linear span of the roots, and let (\cdot, \cdot) be an inner product on $\mathfrak{h}_{\mathbb{R}}^*$ for which $A = (a_{rs})_{r,s \in I} = ((\alpha_r^\vee, \alpha_s))_{r,s \in I}$ is the Cartan matrix of \mathfrak{g} , where $\alpha^\vee = \frac{2}{(\alpha, \alpha)}\alpha$ for $\alpha \in \Delta$. Let $\mathfrak{h}_{\mathbb{R}} \subseteq \mathfrak{h}$ be the real linear span of the coroots h_α , where $\beta(h_\alpha) = (\beta, \alpha^\vee)$ for $\alpha, \beta \in \Delta$.

We further use the following notation. We write $\{\omega_r \mid r \in I\}$ for the fundamental weights in $\mathfrak{h}_{\mathbb{R}}^*$, so $(\omega_r, \alpha_s^\vee) = \delta_{rs}$. The \mathbb{Z} -lattice spanned by $\{\omega_r\}$ is denoted $P \subseteq \mathfrak{h}_{\mathbb{R}}^*$, and P^+ denotes elements expressed as positive linear combinations of this basis. Similarly, the root lattice spanned by the α_r is denoted Q , and its positive span by Q^+ . We write $\text{Char}_K(F)$ for the monoid of monoid homomorphisms from a commutative (additive) monoid $(F, +)$ to a commutative (multiplicative) monoid (K, \cdot) . For $\varepsilon \in \text{Char}_K(Q^+)$ we will abbreviate $\varepsilon_{\alpha_r} = \varepsilon_r$. The unit element of $\text{Char}_K(Q^+)$ will be denoted $+$, while the element ε such that $\varepsilon_r = 0$ for all r will be denoted 0 .

We use the following notation for q -numbers, where $0 < q < 1$ is fixed for the rest of the paper:

- for $r \in \Phi^+$, $q_r = q^{\frac{(\alpha_r, \alpha_r)}{2}}$,
- for $n \geq 0$, $[n]_r = \frac{q_r^n - q_r^{-n}}{q_r - q_r^{-1}}$,
- for $n \geq 0$, $[n]_r! = \prod_{k=1}^n [k]_r$,
- for $m \geq n \geq 0$, $\begin{bmatrix} m \\ n \end{bmatrix}_r = \frac{[m]_r!}{[n]_r! [m-n]_r!}$,
- for $\alpha \in \mathfrak{h}^*$, we define $q^\alpha \in \text{Char}_{\mathbb{C}}(P)$ by $q^\alpha(\omega) = q^{(\alpha, \omega)}$ (where (\cdot, \cdot) has been \mathbb{C} -linearly extended to \mathfrak{h}).

We will only work with unital algebras defined over \mathbb{C} , and correspondingly all tensor products are algebraic tensor products over \mathbb{C} . By a $*$ -algebra A will be meant an algebra A endowed with an anti-linear, anti-multiplicative involution $*$: $A \rightarrow A$. We further assume that the reader is familiar with the theory of Hopf algebras. A *Hopf $*$ -algebra* (H, Δ) is a Hopf algebra whose underlying algebra H is a $*$ -algebra, and whose comultiplication Δ is a $*$ -homomorphism. This implies that the counit is a $*$ -homomorphism, and that the antipode S is invertible with $S^{-1}(h) = S(h^*)^*$ for all $h \in H$.

Definition 1.1. For $\varepsilon, \eta \in \text{Char}_{\mathbb{R}}(Q^+)$, we define $U_q(\mathfrak{g}; \varepsilon, \eta)$ as the universal unital $*$ -algebra generated by couples of elements X_r^\pm , $r \in I$, as well as elements K_ω , $\omega \in P$, such that for all $r, s \in I$ and $\omega, \chi \in P$, we have K_ω self-adjoint, $(X_r^+)^* = X_r^-$ and

$$\begin{aligned} \text{(K)}^q \quad & K_\omega \text{ is invertible and } K_\chi K_\omega^{-1} = K_{\chi - \omega}, \\ \text{(T)}^q \quad & K_\omega X_r^\pm K_\omega^{-1} = q^{\frac{\pm(\omega, \alpha_r)}{2}} X_r^\pm, \\ \text{(S)}^q \quad & \sum_{k=0}^{1-a_{rs}} (-1)^k \begin{bmatrix} 1 - a_{rs} \\ k \end{bmatrix}_r (X_r^\pm)^k X_s^\pm (X_r^\pm)^{1-a_{rs}-k} = 0 \text{ for } r \neq s, \\ \text{(C)}_{\varepsilon, \eta}^q \quad & [X_r^+, X_s^-] = \delta_{rs} \frac{\varepsilon_r K_{\alpha_r}^2 - \eta_r K_{\alpha_r}^{-2}}{q_r - q_r^{-1}}. \end{aligned}$$

When $\varepsilon = \eta = +$ (i.e. $\varepsilon_r = \eta_r = 1$ for all r), we will denote the underlying algebra as $U_q(\mathfrak{g})$. This is a slight variation, obtained by considering $\frac{1}{2}P$ instead of P , of the simply-connected version of the ordinary quantized universal enveloping algebra of \mathfrak{g} [4, Remark 9.1.3].

The $U_q(\mathfrak{g}; \varepsilon, \varepsilon)$ are quantizations of the $*$ -algebras $U(\mathfrak{g}_\varepsilon)$, introduced in Appendix B. Note also that the algebras $U_q(\mathfrak{g}; 0, +)$ are well-known to algebraists, see [20, Section 3] and [17].

Remark that the algebras $U_q(\mathfrak{g}; \varepsilon, \eta)$ with $\prod_r \varepsilon_r \eta_r \neq 0$ are mutually isomorphic as unital algebras. Indeed, as A is invertible, we can choose $b \in \text{Char}_{\mathbb{C}}(P)$ such that $b_{\alpha_r}^4 = \eta_r / \varepsilon_r$ for all $r \in \Phi^+$. We can also choose $a \in \text{Char}_{\mathbb{C}}(Q^+)$ such that a_r is a square root of $b_{\alpha_r}^2 \varepsilon_r = \frac{\eta_r}{b_{\alpha_r}^2}$. Then

$$\phi: U_q(\mathfrak{g}; \varepsilon, \eta) \rightarrow U_q(\mathfrak{g}), \quad \begin{cases} X_r^+ \mapsto a_r X_r^+, \\ X_r^- \mapsto a_r X_r^-, \\ K_\omega \mapsto b_\omega K_\omega \end{cases} \quad (1.1)$$

is a unital isomorphism. However, unless $\text{sgn}(\varepsilon_r \eta_r) = +$ for all r , in which case we can choose $b \in \text{Char}_{\mathbb{R}^+}(P)$ and $a \in \text{Char}_{\mathbb{R}^+}(Q^+)$, this rescaling will not respect the $*$ -structure.

We list some properties of the $U_q(\mathfrak{g}; \varepsilon, \eta)$. The proofs do not differ from those for the well-known case $U_q(\mathfrak{g})$, cf. [31] and [16, Section 4].

Proposition 1.2.

1. Let $U_q(\mathfrak{n}^\pm)$ be the unital subalgebra generated by the X_r^\pm inside $U_q(\mathfrak{g}; \varepsilon, \eta)$. Then $U_q(\mathfrak{n}^\pm)$ is the universal algebra generated by elements X_r^\pm satisfying the relations (S)^q, and in particular does not depend on ε or η .
2. Similarly, let $U_q(\mathfrak{b}^\pm)$ be the subalgebra generated by $U_q(\mathfrak{n}^\pm)$ and $U(\mathfrak{h})$, where $U(\mathfrak{h})$ is the algebra generated by all K_ω . Then $U_q(\mathfrak{b}^\pm)$ is universal with respect to the relations (K)^q, (T)^q and (S)^q.
3. (Triangular decomposition) The multiplication map gives an isomorphism

$$U_q(\mathfrak{n}^+) \otimes U(\mathfrak{h}) \otimes U_q(\mathfrak{n}^-) \rightarrow U_q(\mathfrak{g}; \varepsilon, \eta). \quad (1.2)$$

In particular, the $U_q(\mathfrak{g}; \varepsilon, \eta)$ are non-trivial. Note that the above proposition allows one to identify $U_q(\mathfrak{g}; \varepsilon, \eta)$ and $U_q(\mathfrak{g})$ as *vector spaces*,

$$i_{\varepsilon, \eta}: U_q(\mathfrak{g}; \varepsilon, \eta) \rightarrow U_q(\mathfrak{g}), \quad x_+ x_0 x_- \mapsto x_+ x_0 x_-, \quad x_\pm \in U_q(\mathfrak{n}^\pm), \quad x_0 \in U(\mathfrak{h}).$$

Hence $U_q(\mathfrak{g}; \varepsilon, \eta)$ may also be viewed as $U_q(\mathfrak{g})$ with a deformed product $m_{\varepsilon, \eta}$. Indeed, arguing as in [21, Section 4], one can show that

$$m_{\varepsilon, \eta}(x \otimes y) = \omega_\varepsilon(x_{(1)}, y_{(1)}) x_{(2)} y_{(2)} \omega_\eta(x_{(3)}, y_{(3)})$$

for certain cocycles ω_ε on $U_q(\mathfrak{g})$.

Also the following proposition is immediate.

Proposition 1.3. *For each $\varepsilon, \mu, \eta \in \text{Char}_{\mathbb{R}}(Q^+)$, there exists a unique unital *-homomorphism*

$$\Delta_{\varepsilon, \eta}^\mu: U_q(\mathfrak{g}; \varepsilon, \eta) \rightarrow U_q(\mathfrak{g}; \varepsilon, \mu) \otimes U_q(\mathfrak{g}; \mu, \eta)$$

such that $\Delta_{\varepsilon, \eta}^\mu(K_\omega) = K_\omega \otimes K_\omega$ for all $\omega \in P$ and

$$\Delta_{\varepsilon, \eta}^\mu(X_r^\pm) = X_r^\pm \otimes K_{\alpha_r} + K_{\alpha_r}^{-1} \otimes X_r^\pm, \quad \forall r \in \Phi^+.$$

Proof. From Proposition 1.2 and the case $\varepsilon = \eta = +$, we know that $\Delta_{\varepsilon, \eta}^\mu$ respects the relations (K)^q, (T)^q and (S)^q. The compatibility with the relations (C)^{q, \eta} is verified directly. ■

Lemma 1.4. *The collection $(\{U_q(\mathfrak{g}; \varepsilon, \eta)\}, \{\Delta_{\varepsilon, \eta}^\mu\})$ forms a connected cogroupoid over the index set $\text{Char}_{\mathbb{R}}(Q^+) \cong \mathbb{R}^l$ (cf. Appendix A).*

Proof. One immediately checks the coassociativity condition on the generators. One further can define uniquely a unital *-homomorphism

$$\epsilon_\varepsilon: U_q(\mathfrak{g}; \varepsilon, \varepsilon) \rightarrow \mathbb{C}, \quad \begin{cases} X_r^\pm \mapsto 0, \\ K_\omega \mapsto 1 \end{cases}$$

and unital anti-homomorphism

$$S_{\varepsilon, \eta}: U_q(\mathfrak{g}; \varepsilon, \eta) \rightarrow U_q(\mathfrak{g}; \eta, \varepsilon), \quad \begin{cases} X_r^\pm \mapsto -q^{\pm 1} X_r^\pm, \\ K_\omega \mapsto K_\omega^{-1}. \end{cases}$$

Again, one verifies on generators that these maps satisfy the counit and antipode condition on generators, hence on all elements. ■

We will denote by \triangleleft the associated adjoint action of $U_q(\mathfrak{g}; \eta, \eta)$ on $U_q(\mathfrak{g}; \varepsilon, \eta)$, cf. Appendix A, Definition A.2.

2 Verma module theory

We keep the notation of the previous section.

Definition 2.1. For $\lambda \in \text{Char}_{\mathbb{C}}(P)$, we denote by \mathbb{C}_{λ} the one-dimensional left $U_q(\mathfrak{b}^+)$ -module associated to the character

$$\chi_{\lambda}: U_q(\mathfrak{b}^+) \rightarrow \mathbb{C}, \quad \begin{cases} X_r^+ \mapsto 0, \\ K_{\omega} \mapsto \lambda_{\omega}. \end{cases}$$

We denote

$$M_{\lambda}^{\varepsilon, \eta} = U_q(\mathfrak{g}; \varepsilon, \eta) \otimes_{U_q(\mathfrak{b}^+)} \mathbb{C}_{\lambda}.$$

We denote by $V_{\lambda}^{\varepsilon, \eta}$ the simple quotient of $M_{\lambda}^{\varepsilon, \eta}$ by its maximal non-trivial submodule. Note that such a maximal submodule exists by the triangular decomposition (1.2), as then any non-trivial submodule is a sum of weight spaces with weights distinct from λ . We denote by $v_{\lambda}^{\varepsilon, \eta}$ the highest weight vector $1 \otimes 1$ in either $M_{\lambda}^{\varepsilon, \eta}$ or $V_{\lambda}^{\varepsilon, \eta}$.

For $\lambda \in \text{Char}_{\mathbb{R}}(P)$, we can introduce on $V_{\lambda}^{\varepsilon, \eta}$ a non-degenerate Hermitian form $\langle \cdot, \cdot \rangle$ such that $\langle xv, w \rangle = \langle v, x^*w \rangle$ for all $x \in U_q(\mathfrak{g}; \varepsilon, \eta)$ and $v, w \in V_{\lambda}^{\varepsilon, \eta}$. We will call a form satisfying this property *invariant*. Such a form is then unique up to a scalar. The construction of this form is the same as in the case of $U_q(\mathfrak{g})$ [16, Section 5], [32, Section 2.1.5]. Namely, let $U_q(\mathfrak{n}^{\pm})_{+} \subseteq U_q(\mathfrak{n}^{\pm})$ be the kernel of the restriction of the counit on $U_q(\mathfrak{b}^{\pm})$, and consider the orthogonal decomposition

$$U_q(\mathfrak{g}; \varepsilon, \eta) = U(\mathfrak{h}) \oplus (U_q(\mathfrak{n}^{-})_{+} U_q(\mathfrak{g}; \varepsilon, \eta) + U_q(\mathfrak{g}; \varepsilon, \eta) U_q(\mathfrak{n}^{+})_{+}). \quad (2.1)$$

Let τ denote the projection onto the first summand. Then one first observes that one has a Hermitian form on $M_{\lambda}^{\varepsilon, \eta}$ by defining

$$\langle xv_{\lambda}^{\varepsilon, \eta}, yv_{\lambda}^{\varepsilon, \eta} \rangle = \chi_{\lambda}(\tau(x^*y)), \quad x, y \in U_q(\mathfrak{g}).$$

This is clearly a well-defined invariant form. It necessarily descends to a non-degenerate Hermitian form on $V_{\lambda}^{\varepsilon, \eta}$.

The goal is to find necessary and sufficient conditions for this Hermitian form to be positive-definite, in which case the module is called *unitarizable*. This is in general a hard problem. In the following, we present some partial results, restricting to the case $\eta = +$. We always assume that λ is real-valued, unless otherwise mentioned, and that $M_{\lambda}^{\varepsilon, \eta}$ and $V_{\lambda}^{\varepsilon, \eta}$ have been equipped with the above canonical Hermitian form.

We first consider the case $\varepsilon = \eta = +$, for which the following result is well-known.

Lemma 2.2. *Let $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$. Then the following are equivalent.*

1. *The module $V_{\lambda}^{+,+}$ is unitarizable.*
2. *For all $r \in I$, $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}}$.*
3. *$V_{\lambda}^{+,+}$ is finite-dimensional.*

Proof. For the implications (2) \Rightarrow (3) \Rightarrow (1), see e.g. [4, Corollary 10.1.15, Proposition 10.1.21]. Assume now that $V_{\lambda}^{+,+}$ is unitarizable. Then $V_{\lambda_r}^{+,+}$ is a unitarizable representation of $U_{q_r}(\mathfrak{su}(2))$. By a simple computation using the commutation between the X_r^{\pm} (cf. [9]), we have that

$$\langle (X_r^+)^k (X_r^-)^k v_{\lambda_r}^{+,+}, v_{\lambda_r}^{+,+} \rangle = \prod_{l=1}^k \frac{(q_r^l - q_r^{-l})(q_r^{-l+1} \lambda_{\alpha_r}^2 - q_r^{l-1} \lambda_{\alpha_r}^{-2})}{(q_r - q_r^{-1})^2}.$$

Hence, by unitarity and the fact that $0 < q_r < 1$, we find that $\prod_{l=1}^k (q_r^{l-1} \lambda_{\alpha_r}^{-2} - q_r^{-l+1} \lambda_{\alpha_r}^2) \geq 0$ for all $k \geq 0$. This is only possible if eventually one of the factors becomes zero, i.e. when $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}}$. ■

By a limit argument, we now extend this result to the case $\varepsilon \in \text{Char}_{\{0,1\}}(Q^+)$.

Proposition 2.3. *Suppose $\varepsilon \in \text{Char}_{\{0,1\}}(Q^+)$, and let $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$. Then $V_\lambda^{\varepsilon,+}$ is unitarizable if and only if $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}}$ for all $r \in I$ with $\varepsilon_r = 1$.*

Proof. The ‘only if’ part of the proposition is obvious, since the K_{α_r}, X_r^\pm with $\varepsilon_r = 1$ generate a quantized enveloping algebra $\widetilde{U}_q(\mathfrak{k}_{\text{ss}})$ in its compact (non-simply connected) form.

To show the opposite direction, consider first general $\varepsilon, \eta \in \text{Char}_{\mathbb{R}}(Q^+)$. For $\alpha \in Q^+$, denote by $U_q(\mathfrak{n}^-)(\alpha)$ the finite-dimensional space of elements $X \in U_q(\mathfrak{n}^-)$ with $K_\omega X K_\omega^{-1} = q^{-\frac{1}{2}(\omega, \alpha)} X$. Identifying $U_q(\mathfrak{n}^-)$ with $M_\lambda^{\varepsilon, \eta}$ in the canonical way by means of $x \mapsto xv_\lambda^{\varepsilon, \eta}$, we may interpret the Hermitian forms on the $M_\lambda^{\varepsilon, \eta}$ as a family of Hermitian forms $\langle \cdot, \cdot \rangle_\lambda^{\varepsilon, \eta}$ on $U_q(\mathfrak{n}^-)$. It is easily seen that these vary continuously with ε, η and λ on each $U_q(\mathfrak{n}^-)(\alpha)$. Assume now that $\eta = +$ and $\varepsilon_r > 0$ for all r . In this case $U_q(\mathfrak{g}; \varepsilon, +)$ is isomorphic to $U_q(\mathfrak{g})$ as a $*$ -algebra by a rescaling of the generators by positive numbers, cf. (1.1). By Lemma 2.2, $\langle \cdot, \cdot \rangle_\lambda^{+,+}$ is positive semi-definite if and only if $V_\lambda^{+,+}$ is finite-dimensional, and the latter happens if and only if $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}}$ for all $r \in I$. Hence we get by the above rescaling that $\langle \cdot, \cdot \rangle_\lambda^{\varepsilon,+}$ is positive semi-definite if and only if $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}} \varepsilon_r^{-1}$ for all r .

Fix now a subset S of the simple positive roots, and put $\varepsilon_r = 1$ for $r \in S$. Assume that $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}}$ when $r \in S$. For $r \notin S$ and $m_r \in \mathbb{N}$, define $\varepsilon_r = q_r^{2m_r} \lambda_{\alpha_r}^{-4}$. From the above, we obtain that $\langle \cdot, \cdot \rangle_\lambda^{\varepsilon,+}$ is positive semi-definite. Taking the limit $m_r \rightarrow \infty$ for $r \notin S$, we deduce that $\langle \cdot, \cdot \rangle_\lambda^{\varepsilon,+}$ is positive semi-definite for $\varepsilon_r = \delta_{r \in S}$. ■

The above $V_\lambda^{\varepsilon,+}$ with pre-Hilbert space structure can also be presented more concretely as generalized Verma modules (from which it will be clear that they are not finite dimensional when $\varepsilon_r = 0$ for some r). We will need some preparations, obtained from modifying arguments in [16]. Note that to pass from the conventions in [16] to ours, the q in [16] has to be replaced by $q^{1/2}$, and the coproduct in [16] is also opposite to ours. However, as [16] gives preference to the left adjoint action, while we work with the right adjoint action, most of our formulas will in fact match.

We start with recalling a basic fact.

Lemma 2.4. *Let $\omega \in P^+$, $\varepsilon \in \text{Char}_{\mathbb{R}}(Q^+)$, and consider $K_\omega^{-4} \in U_q(\mathfrak{g}; \varepsilon, +)$. Then $K_\omega^{-4} \triangleleft U_q(\mathfrak{g})$ is finite-dimensional, where \triangleleft is the adjoint action (cf. Definition A.2).*

Proof. One checks that the arguments of [16, Lemma 6.1, Lemma 6.2, Proposition 6.3, Proposition 6.5] are still valid in our setting. ■

Let $\varepsilon \in \text{Char}_{\mathbb{R}}(Q^+)$. Recall that the map τ denoted the projection onto the first summand in (2.1). Let $\tau_{Z, \varepsilon}$ be the restriction of τ to the center $\mathcal{Z}(U_q(\mathfrak{g}; \varepsilon, +))$ of $U_q(\mathfrak{g}; \varepsilon, +)$. This is an ε -modified Harish-Chandra map. The usual reasoning shows that this is a homomorphism into $U(\mathfrak{h})$.

Lemma 2.5. *The map $\tau_{Z, \varepsilon}$ is a bijection between $\mathcal{Z}(U_q(\mathfrak{g}; \varepsilon, +))$ and the linear span of the set $\left\{ \sum_{w \in W} \varepsilon_{\omega - w\omega} q^{(-2w\omega, \rho)} K_{-4w\omega} \mid \omega \in P^+ \right\}$, where $\rho = \sum_r \omega_r$ is the sum of the fundamental weights and W denotes the Weyl group of Δ with its natural action on \mathfrak{h}^* .*

Proof. Note first that $\omega - w\omega$ is inside Q^+ for all $\omega \in P^+$ and $w \in W$, so that $\varepsilon_{\omega-w\omega}$ is well-defined.

Arguing as in [16, Section 8], we can assign to any $\omega \in P^+$ an element z_ω in $\mathcal{Z}(U_q(\mathfrak{g}; \varepsilon, +))$, uniquely determined up to a non-zero scalar, such that

$$K_\omega^{-4} \triangleleft U_q(\mathfrak{g}) = \mathbb{C}z_\omega + (K_\omega^{-4} \triangleleft U_q(\mathfrak{g})_+),$$

where $U_q(\mathfrak{g})_+$ is the kernel of the counit and where \triangleleft denotes the adjoint action. More concretely, as $K_\omega^{-4} \triangleleft U_q(\mathfrak{g})$ is a finite-dimensional right $U_q(\mathfrak{g})$ -module, it is semi-simple, and we have a projection E of $K_\omega^{-4} \triangleleft U_q(\mathfrak{g})$ onto the space of its invariant elements. We can then take $z_\omega = E(K_\omega^{-4})$, and $z_\omega \in \mathcal{Z}(U_q(\mathfrak{g}; \varepsilon, +))$ by Lemma A.3.

Suppose now first that $\varepsilon_r \neq 0$ for all r , and choose $b \in \text{Char}_{\mathbb{C}}(P)$ such that $b_{\alpha_r}^4 = \varepsilon_r^{-1}$. Consider

$$\Psi_\varepsilon: U_q(\mathfrak{g}; \varepsilon, +) \rightarrow U_q(\mathfrak{g}), \quad \begin{cases} X_r^\pm \mapsto b_{\alpha_r}^{-1} X_r^\pm, \\ K_\beta \mapsto b_\beta K_\beta, \end{cases} \quad \beta \in P.$$

This is a unital \triangleleft -equivariant algebra isomorphism, and $\tau_{Z, \varepsilon} = \Psi_\varepsilon^{-1} \circ \tau_{Z, +} \circ \Psi_\varepsilon$. Hence by \triangleleft -equivariance, we find from the computations in [16, Section 8] that, for a non-zero ε -independent scalar c_ω ,

$$\tau_{Z, \varepsilon}(z_\omega) = c_\omega \sum_{\nu \in P^+} \dim((V_\omega)_\nu) \left(\sum_{w \in W} b_{\omega-w\nu}^{-4} q^{-2(w\nu, \rho)} K_{-4w\nu} \right),$$

where $(V_\omega)_\nu$ denotes the weight space at $q^{\frac{1}{2}\nu}$ (i.e. the space of vectors on which the K_ω act as $q^{\frac{1}{2}(\nu, \omega)}$) of the finite-dimensional $U_q(\mathfrak{g})$ -module V_ω with highest weight $q^{\frac{1}{2}\omega}$. But clearly we then only have to sum over those ν with $\omega - \nu \in Q^+$, so that $b_{\omega-w\nu}^{-4} = \varepsilon_{\omega-w\nu}$, and we can write

$$\tau_{Z, \varepsilon}(z_\omega) = c_\omega \sum_{\substack{\nu \in P^+ \\ \omega - \nu \in Q^+}} \dim((V_\omega)_\nu) \left(\sum_{w \in W} \varepsilon_{\omega-w\nu} q^{-2(w\nu, \rho)} K_{-4w\nu} \right). \quad (2.2)$$

Recall now that $U_q(\mathfrak{g}; \varepsilon, \eta)$ can be identified with $U_q(\mathfrak{g})$ as a vector space by a map $i_{\varepsilon, \eta}$. Let us denote by $\triangleleft_{\varepsilon, \eta}$ the image of \triangleleft under this map. Then for $x, y \in U_q(\mathfrak{g})$ fixed, it is easily seen from the triangular decomposition that the $x \triangleleft_{\varepsilon, \eta} y$ live in a fixed finite-dimensional subspace of $U_q(\mathfrak{g})$ as the ε, η vary, and the resulting map $(\varepsilon, \eta) \mapsto x \triangleleft_{\varepsilon, \eta} y$ is then continuous. Furthermore, if V is a finite-dimensional right $U_q(\mathfrak{g})$ -module with space of fixed elements V_{triv} , we can find $p \in U_q(\mathfrak{g})$ such that for any $v \in V$, the element vp is the projection of v onto V_{triv} . It follows from the previous paragraph and the above remarks that when $\varepsilon_r \neq 0$ for any r , we have $i_{\varepsilon, +}(z_\omega) = K_\omega^{-4} \triangleleft_{\varepsilon, +} p_\omega$ for some fixed $p_\omega \in U_q(\mathfrak{g})$. By continuity, it then follows that (2.2) in fact holds for arbitrary $\varepsilon \in \text{Char}_{\mathbb{R}}(Q^+)$.

The conclusion of the argument now follows as in [16, Theorem 8.6]. \blacksquare

Let now $S \subseteq \Phi^+$, and let ε extend the characteristic function of S . Let $U_q(\mathfrak{t}_S)$ be the Hopf $*$ -subalgebra of $U_q(\mathfrak{g}; \varepsilon, +)$ generated by the $K_{\omega_r}^\pm$ with $1 \leq r \leq l$ and X_r^\pm with $r \in S$. Let $U_q(\mathfrak{q}_S^+)$ be the Hopf subalgebra of $U_q(\mathfrak{g}; \varepsilon, +)$ generated by $U_q(\mathfrak{t}_S)$ and all X_r^+ with $r \in I$. It is easy to see that $U_q(\mathfrak{q}_S^+)$ can be isomorphically imbedded into $U_q(\mathfrak{g})$. Let V be a finite-dimensional highest weight representation of $U_q(\mathfrak{t}_S)$ associated to a character in $\text{Char}_{\mathbb{R}^+}(P)$. Then we can extend this to a representation of $U_q(\mathfrak{q}_S^+)$ on V [32, Section 2.3.1], and hence we can form

$$\text{Ind}_\varepsilon(V) = U_q(\mathfrak{g}; \varepsilon, +) \otimes_{U_q(\mathfrak{q}_S^+)} V.$$

The following proposition complements Proposition 2.3.

Proposition 2.6. *Let $S \subseteq \Phi^+$, and let ε restrict to the characteristic function of S . Let V be an irreducible highest weight representation of $U_q(\mathfrak{t}_S)$ associated to a character $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$. Then the $U_q(\mathfrak{g}; \varepsilon, +)$ -representation $\text{Ind}_\varepsilon(V)$ is irreducible.*

Proof. By its universal property, $\text{Ind}_\varepsilon(V)$ can be identified with a quotient of $M_\lambda^{\varepsilon,+}$. We want to show that this quotient coincides with $V_\lambda^{\varepsilon,+}$.

Suppose that $v_{\lambda'}$ is a highest weight vector inside $M_\lambda^{\varepsilon,+}$ at weight λ' different from λ . By Lemma 2.5, it follows that

$$\sum_{w \in W} \varepsilon_{\omega - w\omega} q^{(-2w\omega, \rho)} \lambda'_{-4w\omega} = \sum_{w \in W} \varepsilon_{\omega - w\omega} q^{(-2w\omega, \rho)} \lambda_{-4w\omega}$$

for all $\omega \in P^+$. Now if $w = s_{\alpha_{i_1}} \cdots s_{\alpha_{i_p}}$ in reduced form, with s_α the reflection across the root α , we have

$$\omega - w\omega = \sum_{t=1}^p s_{\alpha_{i_1}} \cdots s_{\alpha_{i_{t-1}}} (\omega - s_{\alpha_{i_t}} \omega),$$

where each term is positive. It follows that we have

$$\sum_{w \in W_S} q^{(-2w\omega, \rho)} \lambda'_{-4w\omega} = \sum_{w \in W_S} q^{(-2w\omega, \rho)} \lambda_{-4w\omega}$$

for all *strictly* dominant ω , where W_S is the Coxeter group generated by reflections around simple roots α_s with $s \in S$.

Taking $\omega = \rho + \omega_r$ with $r \notin S$, we get $(\lambda'_{-4\omega_r} - \lambda_{-4\omega_r})C = 0$ with

$$C = \sum_{w \in W_S} q^{(-2w\rho, \rho)} \lambda_{-4w\rho} > 0.$$

Hence $\lambda'_{\omega_r} = \lambda_{\omega_r}$ for all $r \notin S$. We deduce that $v_{\lambda'} \in U_q(\mathfrak{t}_S)v_\lambda$, and so the image of $v_{\lambda'}$ in $\text{Ind}_\varepsilon(V)$ is zero. This implies that $\text{Ind}_\varepsilon(V) = V_\lambda^{\varepsilon,+}$. \blacksquare

The case $\varepsilon \in \text{Char}_{\{-1,0,1\}}(Q^+)$ is not so easy to treat in general. In the following, we will restrict ourselves to the case where we have one $\varepsilon_t < 0$ at a root satisfying a particular condition, while $\varepsilon_r \geq 0$ for $r \neq t$. This will correspond precisely to the ‘Hermitian symmetric’ case.

Theorem 2.7. *Let ε be such that there is a unique simple root α_t with $\varepsilon_t < 0$, while $\varepsilon_r \in \{0, 1\}$ for $r \neq t$. Assume moreover that α_t appears with multiplicity at most 1 in each positive root. Let $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$. Then $V_\lambda^{\varepsilon,+}$ is unitarizable if and only if $\lambda_{\alpha_r}^4 \in q_r^{2\mathbb{N}}$ for all r with $\varepsilon_r \neq 1$.*

One can indeed check by a case-by-case analysis, using for example the tables in [23, Appendix C], that the Vogan diagram associated to a sign pattern will determine a Hermitian symmetric space precisely when the above multiplicity condition is satisfied.

Proof. By a same kind of limiting argument as in Proposition 2.3, the general case can be deduced from the case with $\varepsilon_r = 1$ for $r \neq t$.

Suppose then that $\varepsilon_t < 0$ and $\varepsilon_r = 1$ for $r \neq t$, where α_r appears with multiplicity at most 1 in each positive root. Write $S = \Phi^+ \setminus \{t\}$. We have the algebra automorphism $\phi: U_q(\mathfrak{g}; \varepsilon, +) \rightarrow U_q(\mathfrak{g})$ appearing in (1.1). By means of this isomorphism, we obtain a natural isomorphism $M_\lambda^{\varepsilon,+} \cong M_\gamma^{+,+}$, where $\gamma \in \text{Char}_{\mathbb{C}}(P)$ is such that $\gamma_{\alpha_r}^4 = \varepsilon_r \lambda_{\alpha_r}^4$ for all r . In particular, $\gamma_{\alpha_t}^2 \in \mathbb{C} \setminus \mathbb{R}$. By the condition we assume on α_r , we can apply [18, Proposition 5.13] and deduce that $\text{Ind}_\varepsilon(V_\lambda)$ is irreducible, where V_λ denotes the irreducible representation of $U_q(\mathfrak{q}_S^+)$ at highest weight λ . Hence the signatures of the Hermitian inner products on the $\text{Ind}_\varepsilon(V_\lambda)$ are constant as $\varepsilon_t < 0$

varies. Indeed, these spaces can be identified canonically with a fixed quotient of $U_q(\mathfrak{n}^-)$, see [32, Proposition 2.81], and then the Hermitian inner products clearly form a continuous family as ε varies.

From Proposition 2.6, we know that the Hermitian inner product on $\text{Ind}_\varepsilon(V_\lambda)$ for $\varepsilon_t = 0$ is positive definite. It follows that the Hermitian inner product on a weight space of $\text{Ind}_\varepsilon(V_\lambda)$ is positive for $\varepsilon_t < 0$ small. As the signature is constant, this holds for all $\varepsilon_t < 0$. ■

3 Quantized homogeneous spaces

Definition 3.1. For $\varepsilon, \eta \in \text{Char}_{\mathbb{R}}(Q^+)$, we denote by $U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}$ the space of locally finite vectors in $U_q(\mathfrak{g}; \varepsilon, \eta)$ with respect to the right adjoint action by $U_q(\mathfrak{g}; \eta, \eta)$ (cf. Definition A.2),

$$U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}} = \{x \in U_q(\mathfrak{g}; \varepsilon, \eta) \mid \dim(x \triangleleft U_q(\mathfrak{g}; \eta, \eta)) < \infty\}.$$

It is easily seen that the space $U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}$ is a $*$ -subalgebra of $U_q(\mathfrak{g}; \varepsilon, \eta)$ (cf. [16, Corollary 2.3]), and in the following it will always be treated as a right $U_q(\mathfrak{g})$ -module by \triangleleft .

A similar definition of ${}_{\text{fin}}U_q(\mathfrak{g}; \varepsilon, \eta)$ can be made with respect to the left adjoint action of $U_q(\mathfrak{g}; \varepsilon, \varepsilon)$, and the two resulting algebras $U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}$ and ${}_{\text{fin}}U_q(\mathfrak{g}; \varepsilon, \eta)$ should in some sense be seen as dual to each other. For example, the $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ will lead to compact quantum homogeneous spaces, while the ${}_{\text{fin}}U_q(\mathfrak{g}; \varepsilon, +)$ should lead to non-compact quantum homogeneous spaces such as quantum bounded symmetric domains [32]. However, in this paper we will restrict ourselves to the compact case.

The $U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}$ are sufficiently large, as the next proposition shows, extending Lemma 2.4.

Proposition 3.2. *As a right $U_q(\mathfrak{g})$ -module, $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ is generated by the K_ω^{-4} with $\omega \in P^+$. The algebra generated by $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ and the $K_{\omega_r}^4$ equals the subalgebra of $U_q(\mathfrak{g}; \varepsilon, +)$ generated by the $K_{\omega_r}^{\pm 4}$ and the $K_{\alpha_r} X_r^\pm$.*

Proof. Again, the proof of [16, Theorem 6.4] can be directly modified. ■

Note that for $\varepsilon_r \neq 0$ for all r , the above proposition follows more straightforwardly from [16] by a rescaling argument.

The dependence of $U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}$ on ε and η is weaker than for $U_q(\mathfrak{g}; \varepsilon, \eta)$ itself. We consider a special case in the following lemma. Recall that A denotes the Cartan matrix.

Lemma 3.3. *Consider $\varepsilon, \eta \in \text{Char}_{\mathbb{R} \setminus \{0\}}(Q^+)$, and write $\text{sgn}(\varepsilon_r/\eta_r) = (-1)^{x_r}$. If χ is in the range of $A \bmod 2$, then $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}} \cong U_q(\mathfrak{g}; \eta, +)_{\text{fin}}$ as right $U_q(\mathfrak{g})$ -module $*$ -algebras.*

Proof. Choose $b \in \text{Char}_{\mathbb{C}}(P)$ such that $b_{\alpha_r}^4 = \eta_r/\varepsilon_r$. Then

$$\Psi_{\varepsilon, \eta}: U_q(\mathfrak{g}; \varepsilon, +) \rightarrow U_q(\mathfrak{g}; \eta, +), \quad \begin{cases} X_r^\pm \mapsto b_{\alpha_r}^{-1} X_r^\pm, \\ K_\omega \mapsto b_\omega K_\omega \end{cases}$$

is a unital \triangleleft -equivariant algebra isomorphism. Hence $\Psi_{\varepsilon, \eta}$ induces a unital \triangleleft -equivariant algebra isomorphism $\psi_{\varepsilon, \eta}: U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}} \rightarrow U_q(\mathfrak{g}; \eta, +)_{\text{fin}}$.

As the K_ω^{-4} with $\omega \in P^+$ generate $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ as a module, $\psi_{\varepsilon, \eta}$ will be $*$ -preserving if and only if $b_{\alpha_r}^4 \in \mathbb{R}$ for all r . This can be realized if we can find $c_r \in \{-1, 1\}$ such that $\prod_s c_s^{a_{sr}} = \text{sgn}(\eta_r/\varepsilon_r)$, which is equivalent with the condition appearing in the statement of the lemma. ■

In particular, we find for example that $U_q(\mathfrak{sl}(2m+1); \varepsilon, +)_{\text{fin}}$ for $m \in \mathbb{N}_0$ is independent of the choice of $\varepsilon \in \text{Char}_{\mathbb{R} \setminus \{0\}}(Q^+)$. On the other hand, $U_q(\mathfrak{sl}(2); \varepsilon, +)_{\text{fin}}$ are mutually non-isomorphic as $*$ -algebras for $\varepsilon \in \{-1, 0, 1\}$, see [8].

Definition 3.4. Let $\varepsilon, \eta \in \text{Char}_{\mathbb{R}}(Q^+)$, $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$, and let $V_{\lambda}^{\varepsilon, \eta}$ be the irreducible highest weight module of $U_q(\mathfrak{g}; \varepsilon, \eta)$ at λ with associated representation $\pi_{\lambda}^{\varepsilon, \eta}$. We write

$$B_{\lambda}(\mathfrak{g}; \varepsilon, \eta) = \pi_{\lambda}^{\varepsilon, \eta}(U_q(\mathfrak{g}; \varepsilon, \eta))$$

and

$$B_{\lambda}^{\text{fin}}(\mathfrak{g}; \varepsilon, \eta) = \pi_{\lambda}^{\varepsilon, \eta}(U_q(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}).$$

Remark 3.5. The space $B_{\lambda}^{\text{fin}}(\mathfrak{g}; \varepsilon, \eta)$ is not defined as the space $B_{\lambda}(\mathfrak{g}; \varepsilon, \eta)_{\text{fin}}$ of locally finite \triangleleft -elements in $B_{\lambda}(\mathfrak{g}; \varepsilon, \eta)$, although conceivably they are the same in many cases. In the case $q = 1$, the equality of these two algebras goes by the name of the Kostant problem, cf. [19, Remark 3].

Notation 3.6. We will use the following notation for particular elements in the $B_{\lambda}(\mathfrak{g}; \varepsilon, +)$:

$$\begin{aligned} Z_r &= \pi_{\lambda}^{\varepsilon, +}(K_{\omega_r}^{-4}), & X_r &= q_r^{1/2}(q_r^{-1} - q_r)\pi_{\lambda}^{\varepsilon, +}(K_{\alpha_r}K_{\omega_r}^{-4}X_r^+), \\ Y_r &= X_r^*, & W_r &= \pi_{\lambda}^{\varepsilon, +}(K_{\alpha_r}^4K_{\omega_r}^{-8}), \\ T_r &= (q_r - q_r^{-1})^2\pi_{\lambda}^{\varepsilon, +}(K_{\alpha_r}^2K_{\omega_r}^{-4}X_r^+X_r^-) + \varepsilon_r q_r^{-1}\pi_{\lambda}^{\varepsilon, +}(K_{\alpha_r}^4K_{\omega_r}^{-4}) + q_r\pi_{\lambda}^{\varepsilon, +}(K_{\omega_r}^{-4}) \\ &= (q_r - q_r^{-1})^2\pi_{\lambda}^{\varepsilon, +}(K_{\alpha_r}^2K_{\omega_r}^{-4}X_r^-X_r^+) + \varepsilon_r q_r\pi_{\lambda}^{\varepsilon, +}(K_{\alpha_r}^4K_{\omega_r}^{-4}) + q_r^{-1}\pi_{\lambda}^{\varepsilon, +}(K_{\omega_r}^{-4}). \end{aligned}$$

The following commutation relations will be needed later on.

Lemma 3.7. *The elements W_r and T_r commute with X_r, Y_r, Z_r, T_r and W_r . Moreover,*

$$X_r Z_r = q_r^2 Z_r X_r, \quad Y_r Z_r = q_r^{-2} Z_r Y_r$$

and

$$X_r Y_r = -\varepsilon_r W_r + q_r T_r Z_r - q_r^2 Z_r^2, \quad Y_r X_r = -\varepsilon_r W_r + q_r^{-1} T_r Z_r - q_r^{-2} Z_r^2.$$

We further have that T_r and W_r are invariant under $\triangleleft X_r^{\pm}$ and $\triangleleft K_{\omega}$, while

$$\begin{aligned} X_r \triangleleft X_r^+ &= 0, & Y_r \triangleleft X_r^+ &= -q_r^{1/2}(q_r^{-1} + q_r)Z_r + q_r^{1/2}T_r, \\ X_r \triangleleft K_{\omega} &= q^{-\frac{(\omega, \alpha_r)}{2}}X_r, & Y_r \triangleleft K_{\omega} &= q^{\frac{(\omega, \alpha_r)}{2}}X_r, \\ X_r \triangleleft X_r^- &= q_r^{-1/2}(q_r^{-1} + q_r)Z_r - q_r^{-1/2}T_r, & Y_r \triangleleft X_r^- &= 0 \end{aligned}$$

and

$$Z_r \triangleleft X_r^+ = q_r^{1/2}X_r, \quad Z_r \triangleleft K_{\omega} = 0, \quad Z_r \triangleleft X_r^- = -q_r^{-1/2}Y_r.$$

Finally, all elements X_r, Y_r, Z_r, T_r, W_r are inside $B_{\lambda}^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$.

Proof. All these assertions follow from straightforward computations. As the $Z_r = \pi_{\lambda}^{\varepsilon, +}(K_{\omega_r}^{-4})$ and W_r are in $B_{\lambda}^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ by Proposition 3.2, and the latter is \triangleleft -stable, it follows from the above computations that also X_r, Y_r and T_r are in $B_{\lambda}^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$. ■

Proposition 3.8. *The only \triangleleft -invariant elements in $B_{\lambda}^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ are scalar multiples of the unit element.*

Proof. Assume that $x \in U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ with $\pi_{\lambda}^{\varepsilon, +}(x)$ invariant. As $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ is a semi-simple right $U_q(\mathfrak{g})$ -module, we have an equivariant projection E of $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ onto the $*$ -algebra of its invariant elements. The latter is simply the center $\mathcal{Z}(U_q(\mathfrak{g}; \varepsilon, +))$ of $U_q(\mathfrak{g}; \varepsilon, +)$, by Lemma A.3. As $\pi_{\lambda}^{\varepsilon, +}$ is \triangleleft -equivariant by construction, we deduce that $\pi_{\lambda}^{\varepsilon, +}(x) = \pi_{\lambda}^{\varepsilon, +}(E(x))$. But the latter is a scalar. ■

Remark 3.9. An alternative proof consists in applying Schur's lemma to the simple module $V_\lambda^{\varepsilon,+}$. Indeed, $x \in B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ is \triangleleft -invariant if and only if it commutes with all $\pi_\lambda^{\varepsilon,+}(y)$ for $y \in U_q(\mathfrak{g}; \varepsilon, +)$. As $V_\lambda^{\varepsilon,+}$ is simple, Schur's lemma implies that the algebra of \triangleleft -invariant elements in $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ forms a field of countable dimension over \mathbb{C} , hence coincides with \mathbb{C} . (I would like the referee for pointing out this approach).

Proposition 3.10. *Let (V, π) be a $*$ -representation of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ on a pre-Hilbert space. Then π is bounded.*

The proof is based on an argument which is well-known in the setting of compact quantum groups.

Proof. As $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ consists of locally finite elements, any $b \in B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ can be written as a finite linear combination of elements $b_i \in B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ for which there exists a finite-dimensional $*$ -representation π of $U_q(\mathfrak{g})$ on a Hilbert space such that $b_i \triangleleft h = \sum_j \pi_{ij}(h)b_j$ for all $h \in U_q(\mathfrak{g})$, the π_{ij} being the matrix components with respect to some orthogonal basis. An easy computation shows that $\sum_i b_i^* b_i$ is an invariant element, hence a scalar by Proposition 3.8. Hence there exists $C \in \mathbb{R}^+$ such that for any $\xi \in V$ and any i , we have $\|\pi(b_i)\xi\| \leq C\|\xi\|$. We deduce that the element $\pi(b)$ is bounded. ■

Definition 3.11. A $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ -module V is called a *highest weight module* if there exists a cyclic vector $v \in V$ which is annihilated by all X_r and which is an eigenvector for all Z_r with non-zero eigenvalue. A pre-Hilbert space structure on V is called *invariant* if $\langle x\xi, \eta \rangle = \langle \xi, x^*\eta \rangle$ for all $\xi, \eta \in V$ and $x \in B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$.

We aim to show that the $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ have only a finite number of non-equivalent irreducible highest weight modules. Of course, each $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ admits at least the highest weight module $V_\lambda^{\varepsilon,+}$. Also note that, by an easy argument, each highest weight module decomposes into a direct sum of joint weight spaces for the Z_r .

Proposition 3.12. *Each $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ admits only a finite number of non-equivalent irreducible highest weight modules.*

Proof. As the statement does not depend on the $*$ -structure, we may by rescaling restrict to the case that $\varepsilon_r \in \{0, 1\}$ for all r upon allowing $\lambda \in \text{Char}_{\mathbb{C}}(P)$.

By Proposition 3.2 and the fact that any highest weight module is semi-simple for the torus part, it is easily argued that any irreducible highest weight module of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ is obtained by restriction of a $U_q(\mathfrak{g}; \varepsilon, +)$ -module $V_{\lambda'}^{\varepsilon,+}$ for some $\lambda' \in \text{Char}_{\mathbb{C}}(P)$. As the center of $U_q(\mathfrak{g}; \varepsilon, +)$ acts by the same character on $V_\lambda^{\varepsilon,+}$ and $V_{\lambda'}^{\varepsilon,+}$, we find by Lemma 2.5 that the expression $\sum_{w \in W} \varepsilon_{\omega - w\omega} q^{-2(w\omega, \rho)} \lambda_{-4w\omega}$ remains the same upon replacing λ by λ' , for each $\omega \in P^+$. Writing S for the set of r with $\varepsilon_r = 0$, it follows as in the proof of Lemma 2.5 that

$$\sum_{w \in W_S} q^{-2(w\omega, \rho)} \lambda_{-4w\omega} = \sum_{w \in W_S} q^{-2(w\omega, \rho)} \lambda'_{-4w\omega}$$

for all $\omega \in P^{++}$, the strictly dominant weights. As (invertible) characters on a commutative semi-group are linearly independent, and as $P^{++} - P^{++} = P$, it follows that the functions $\omega \rightarrow q^{-2(w\omega, \rho)} \lambda_{-4w\omega}$ and $\omega \rightarrow q^{-2(w\omega, \rho)} \lambda'_{-4w\omega}$ on P lie in the same W_S -orbit. As the highest weight vector in an irreducible highest weight module is uniquely determined up to a scalar, and as the equivalence classes of such highest weight modules are then determined by the associated eigenvalue of the $Z_r = \pi_\lambda^{\varepsilon,+}(K_{\omega_r}^{-4})$, this is sufficient to prove the proposition. ■

Remark that the above proof also gives the upper bound $|W_S|$ for the number of inequivalent highest weight representations, but of course this estimate is not sharp if one only considers unitarizable representations.

Proposition 3.13. *Let π be a $*$ -representation of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ on a Hilbert space \mathcal{H} . If 0 is not in the point-spectrum of any of the Z_r , then \mathcal{H} is a (possibly infinite) direct sum of completions (\mathcal{H}_k, π_k) of unitarizable highest weight modules of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$.*

Proof. By a direct integral decomposition, and using Proposition 3.12, it is sufficient to show that any such *irreducible* $*$ -representation π of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ on a Hilbert space \mathcal{H} is the completion of a highest weight module with invariant pre-Hilbert space structure. We then argue as in [28, Section 3]. Write χ_X for the characteristic function of a set. By assumption, there exists $t \in \mathbb{R}^l$ with $t_r \neq 0$ for all r and $P_t = \chi_{\prod_r [q_r t_r, t_r]}(\pi(Z_1), \dots, \pi(Z_l))$ non-zero. Suppose now that r is such that $\pi(X_r)P_t \neq 0$. From the commutation relations between the X_r and the Z_s , we deduce that $P_{(t_1, \dots, q_r^{-2} t_r, \dots, t_l)} \neq 0$. As the $\pi(Z_r)$ are bounded, this process must necessarily stop. Hence we may choose t such that $P_t \neq 0$ but $\pi(X_r)P_t = 0$ for all r .

Let V be the union of the images of the spectral projections of (Z_1, \dots, Z_l) corresponding to the $\prod_r (\mathbb{R} \setminus (-\frac{1}{n}, \frac{1}{n}))$ with $n \in \mathbb{N}$. As $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ is spanned by elements which skew-commute with the Z_r , it follows that V is a $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ -module on which the $\pi(Z_r)$ are invertible linear maps. This entails that the restriction of π to V can be extended to a representation $\tilde{\pi}$ of $B^{\text{fin}}(\mathfrak{g}; \varepsilon, +)_{\text{ext}}$, the sub $*$ -algebra of $B(\mathfrak{g}; \varepsilon, +)$ generated by X_r, Y_r and the $Z_r^{\pm 1}$ (which contains $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ by Proposition 3.2). Note that this $*$ -algebra admits a triangular decomposition (in the obvious way with respect to the above generators).

Pick now a non-zero $\xi \in P_t \mathcal{H}$. Suppose that ξ were not in the pure point spectrum of some $\pi(Z_r)$. Then we can find $q_r t_r < a < t_r$ such that $\chi_{[q_r t_r, a]}(\pi(Z_r))\xi \neq 0 \neq \chi_{(a, t_r]}(\pi(Z_r))\xi$. However, $[q_1 t_1, t_1] \times \dots \times [q_r t_r, a] \times \dots \times [q_l t_l, t_l] \cap \prod_s [q_s^{2k_s+1} t_s, q_s^{2k_s} t_s] = \emptyset$ for all $k_s \in \mathbb{N}$ with at least one $k_s > 0$. From the commutation relations between the Y_s and $Z_{s'}$, and the fact that $\pi(X_s)\xi = 0$ for all s , we deduce that $\chi_{[q_r t_r, a]}(\pi(Z_r))\xi$ is orthogonal to the $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)_{\text{ext}}$ -module spanned by $\chi_{(a, t_r]}(\pi(Z_r))\xi$. As π is irreducible, this would entail $\chi_{[q_r t_r, a]}(\pi(Z_r))\xi = 0$. Having arrived at a contradiction, we conclude that ξ is a joint eigenvector of all $\pi(Z_r)$.

As ξ is annihilated by all $\pi(X_r)$ and is a joint eigenvector of all $\pi(Z_r)$, the module generated by it is a highest weight module. As π was irreducible, this module must necessarily be dense in \mathcal{H} , and the proposition is proven. \blacksquare

We now want to consider analytic versions of the $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$.

Definition 3.14. Let B be a unital $*$ -algebra. We say that B admits a *universal C^* -envelope* if there exists a non-trivial unital C^* -algebra C together with a unital $*$ -homomorphism $\pi_u: B \rightarrow C$ of unital $*$ -algebras such that any $*$ -homomorphism $B \rightarrow D$ with D a unital C^* -algebra factors through C .

Of course, the above C^* -algebra C is then uniquely determined up to isomorphism.

Definition 3.15. We define $\text{Pol}(\mathbb{G}_+^q)$ to be the Hopf $*$ -algebra inside the dual of $U_q(\mathfrak{g})$ which is spanned by the matrix coefficients of finite-dimensional highest weight representations of $U_q(\mathfrak{g})$ associated to positive characters. We define

$$\alpha_\lambda^{\varepsilon, +}: B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +) \rightarrow \text{Pol}(\mathbb{G}_+^q) \otimes B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$$

as the comodule $*$ -algebra structure dual to the module $*$ -algebra structure \triangleleft by $U_q(\mathfrak{g})$.

Note that the latter definition makes sense, since $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ is integrable as a right $U_q(\mathfrak{g})$ -module.

It is known [27] that $\text{Pol}(\mathbb{G}_+^q)$ admits a universal C^* -algebraic envelope $C(\mathbb{G}_+^q)$, which becomes a compact quantum group in the sense of [33]. We will denote by $\varphi_{\mathbb{G}_+^q}$ the invariant state on $C(\mathbb{G}_+^q)$, which is faithful by co-amenability of \mathbb{G}_+^q .

Lemma 3.16. *Assume that $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ admits at least one $*$ -representation on a Hilbert space. Then $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ admits a C^* -algebraic envelope $C_\lambda(\mathfrak{g}; \varepsilon, +)$.*

Proof. The universal C^* -algebraic envelope of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ exists for precisely the same reason as in Proposition 3.10, since for any element $b \in B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ there exists a universal constant C_b such that $\|\pi(b)\| \leq C$ for all $*$ -representations π of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ by bounded operators on a Hilbert space. As $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ admits at least one $*$ -representation, we have $C_\lambda(\mathfrak{g}; \varepsilon, +) \neq 0$. ■

Remark 3.17. If $V_\lambda^{\varepsilon,+}$ is unitarizable, it is of course clear that we get a *faithful* map from $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ into $C_\lambda(\mathfrak{g}; \varepsilon, +)$.

Lemma 3.18. *Let $C_\lambda(\mathfrak{g}; \varepsilon, +)$ be the universal C^* -envelope of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$, whenever it exists. Then $\alpha_\lambda^{\varepsilon,+}$ induces a C^* -algebraic coaction by $C(\mathbb{G}_+^q)$ on $C_\lambda(\mathfrak{g}; \varepsilon, +)$.*

Proof. The map $\alpha_\lambda^{\varepsilon,+}$ gives a C^* -representation of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ into $C(\mathbb{G}_+^q) \otimes C_\lambda(\mathfrak{g}; \varepsilon, +)$, which hence factors over $C_\lambda(\mathfrak{g}; \varepsilon, +)$. It is straightforward to argue that this is a C^* -algebraic coaction. ■

As $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ only had scalar multiples of the unit as invariants, it follows that the coaction $\alpha_\lambda^{\varepsilon,+}$ on $C_\lambda(\mathfrak{g}; \varepsilon, +)$ is *ergodic* [3], i.e. if $\alpha_\lambda^{\varepsilon,+}(x) = 1 \otimes x$, then $x \in \mathbb{C}1$. We will write $\varphi_\lambda^{\varepsilon,+}$ for the unique invariant state on $C_\lambda(\mathfrak{g}; \varepsilon, +)$, so

$$(\varphi_{\mathbb{G}_+^q} \otimes \iota)\alpha_\lambda^{\varepsilon,+}(x) = \varphi_\lambda^{\varepsilon,+}(x)1, \quad \forall x \in C_\lambda(\mathfrak{g}; \varepsilon, +).$$

As $\varphi_{\mathbb{G}_+^q}$ is faithful, also $\varphi_\lambda^{\varepsilon,+}$ is faithful.

Notation 3.19. We write $\theta_{\lambda, \text{reg}}^{\varepsilon,+}$ for the GNS-representation of $(C_\lambda(\mathfrak{g}; \varepsilon, +), \varphi_\lambda^{\varepsilon,+})$, and $W_\lambda(\mathfrak{g}; \varepsilon, +)$ for the von Neumann algebraic completion of $C_\lambda(\mathfrak{g}; \varepsilon, +)$ in this GNS-representation.

From Lemma 3.16, it follows that a C^* -algebraic envelope of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ exists if $V_\lambda^{\varepsilon,+}$ is unitarizable. In this case, we can say something explicit about $W_\lambda(\mathfrak{g}; \varepsilon, +)$.

Theorem 3.20. *Assume that $V_\lambda^{\varepsilon,+}$ is unitarizable. Then $W_\lambda(\mathfrak{g}; \varepsilon, +)$ is a finite direct sum of type I factors.*

The proof will make use of the following standard lemma.

Lemma 3.21. *Let A be a unital C^* -algebra with faithful state φ . Let M be the von Neumann algebra closure of A in its GNS-representation with respect to φ . Let π be a representation of A on a Hilbert space \mathcal{H} such that there exists a faithful state $\omega \in B(\mathcal{H})_*$ with $\omega \circ \pi = \varphi$. Then π extends to a normal faithful $*$ -representation of M .*

Proof. As ω is faithful, the bicommutant $\pi(A)''$ is faithfully represented on the GNS-space $\mathcal{L}^2(\pi(A), \omega)$. The unitary $U: \mathcal{L}^2(A, \varphi) \rightarrow \mathcal{L}^2(\pi(A), \omega)$ induced by π then provides an isomorphism $M \rightarrow \pi(A)''$ extending π . ■

Proof of Theorem 3.20. As mentioned, the C^* -algebraic envelope $C_\lambda(\mathfrak{g}; \varepsilon, +)$ certainly exists.

To prove the remaining part of the theorem, we first make some preparations. Recall that the dual of $U_q(\mathfrak{sl}(2, \mathbb{C}))$ can be identified with $\text{Pol}(SU_q(2))$, Woronowicz's twisted quantum $SU(2)$ -group [34]. It is well-known that its von Neumann algebraic envelope $\mathcal{L}^\infty(SU_q(2))$ is

isomorphic to $B(l^2(\mathbb{N})) \otimes \mathcal{L}(\mathbb{Z})$, and as such admits a faithful representation on the Hilbert space $\mathcal{H}_+ = l^2(\mathbb{N}) \otimes l^2(\mathbb{Z})$ (cf. [27, 30]).

More generally, write $U_{q_r}(\mathfrak{su}(2))$ for the sub-Hopf- $*$ -algebra generated by the X_r^\pm and $K_{\alpha_r}^{\pm 1}$ inside $U_q(\mathfrak{g})$. By duality, one obtains a surjective $*$ -homomorphism $\gamma_r: \text{Pol}(\mathbb{G}_+^q) \rightarrow \text{Pol}(SU_{q_r}(2))$. This induces a $*$ -representation of $\text{Pol}(\mathbb{G}_+^q)$ on \mathcal{H}_+ , which we will denote by the same symbol γ_r . Suppose now that $t = (r_1, \dots, r_n)$ is an ordered n -tuple of elements in I . Then we obtain a $*$ -representation of $\text{Pol}(G_+^q)$ on $\mathcal{H}_+^{\otimes n}$ by means of the $*$ -homomorphism

$$\gamma_t = (\gamma_{r_1} \otimes \cdots \otimes \gamma_{r_n}) \circ \Delta_{\mathbb{G}_+^q}^{(n)},$$

where $\Delta_{\mathbb{G}_+^q}^{(n)}$ denotes the n -fold coproduct. Let now $t_0 = (r_1, \dots, r_N)$ be such that $w_0 = s_{r_1} \cdots s_{r_N}$ is a reduced expression for the longest element in the Weyl group of \mathfrak{g} . By [30], we know that $\varphi_{\mathbb{G}_+^q}$ can be realized as $\omega \circ \gamma_{t_0}$ for some faithful normal state $\omega \in B(\mathcal{H}_+^{\otimes N})_*$. By Lemma 3.21, this implies that γ_{t_0} can be extended to a faithful normal $*$ -representation of $\mathcal{L}^\infty(G_+^q)$.

Let us turn now to the proof of the theorem. The main step is to prove that the point-spectrum of $\theta_{\lambda, \text{reg}}^{\varepsilon, +} \left(\prod_{r=1}^l Z_r \right)$ does not contain zero. Indeed, if this is the case, then we can use Proposition 3.13 to conclude that $\theta_{\lambda, \text{reg}}^{\varepsilon, +}$ decomposes into a direct integral of highest weight modules for $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$. It then follows that $W_\lambda(\mathfrak{g}; \varepsilon, +)$ is simply $\oplus_{k=1}^m B(\mathcal{H}_k)$ with \mathcal{H}_k the Hilbert space completion of the highest weight modules which appear in $\theta_{\lambda, \text{reg}}^{\varepsilon, +}$.

To show that $\theta_{\lambda, \text{reg}}^{\varepsilon, +} \left(\prod_{r=1}^l Z_r \right)$ does not contain zero in its pointspectrum, it is sufficient to show that the operator $(\gamma_{t_0} \otimes \theta_\lambda^{\varepsilon, +}) \left(\alpha_\lambda^{\varepsilon, +} \left(\prod_{r=1}^l Z_r \right) \right)$ does not contain zero in its pointspectrum, where $\theta_\lambda^{\varepsilon, +}$ is the $*$ -representation of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ on the Hilbert space completion $\mathcal{H}_\lambda^{\varepsilon, +}$ of $V_\lambda^{\varepsilon, +}$. Indeed, the invariant state $\varphi_\lambda^{\varepsilon, +} = (\varphi_{\mathbb{G}_+^q} \otimes \text{id}) \alpha_\lambda^{\varepsilon, +}$ can be extended to a faithful normal functional on $B(\mathcal{H}_\lambda^{\varepsilon, +} \otimes \mathcal{H}_+^{\otimes N})$, which implies, again by Lemma 3.21, that $(\gamma_{t_0} \otimes \theta_\lambda^{\varepsilon, +}) \alpha_\lambda^{\varepsilon, +}$ extends to a faithful normal $*$ -representation of $W_\lambda(\mathfrak{g}; \varepsilon, +)$.

Finally, to show that $(\theta_\lambda^{\varepsilon, +} \otimes \gamma_{t_0}) \left(\alpha_\lambda^{\varepsilon, +} \left(\prod_{r=1}^l Z_r \right) \right)$ does not contain zero in its pointspectrum, we can reason by induction, using the following lemma. ■

Lemma 3.22. *Let (V, π) be an irreducible highest weight module for $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ with an invariant pre-Hilbert space structure, and let \mathcal{H} be the completion of V . Fix $r \in I$, and put $\pi_r = (\pi \otimes \gamma_r) \alpha_\lambda^{\varepsilon, +}$. Then $\pi_r \left(\prod_{s=1}^l Z_s \right)$ does not contain 0 in its point spectrum.*

Proof. It is easy to see that $\pi_r(Z_s) = 1 \otimes \pi(Z_s)$ for $r \neq s$, so that none of these operators have zero in their point-spectrum. Let now A_r be the sub- $*$ -algebra of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ generated by Z_r, T_r, X_r, Y_r, W_r , see Lemma 3.7. By that lemma, A_r is stable under the right action by $U_{q_r}(\mathfrak{su}(2))$, and W_r and T_r are invariants in the center of A_r . By invariance, $\pi_r(W_r) = 1 \otimes \pi(W_r)$ and $\pi_r(T_r) = 1 \otimes \pi(T_r)$ are bounded self-adjoint operators. Hence, to investigate the spectrum of $\pi_r(Z_r)$, we may by disintegration treat the above operators as scalars, say w_r and t_r . Denote the resulting quotient of A_r by $A_r(w_r, t_r)$.

From Lemma 3.7, we find commutation relations between the generators X_r, Y_r and Z_r of $A_r(w_r, t_r)$, as well as the resulting action of $U_{q_r}(\mathfrak{su}(2))$. It follows that $A_r(w_r, t_r)$ is an equivariant quotient of a generalized Podleś sphere $S_{q_r, \tau}^2$ for $SU_{q_r}(2)$ [28], for some τ depending on t_r and w_r . Moreover, as $\varphi_{SU_{q_r}(2)}$ can be realized on the Hilbert space \mathcal{H}_+ , the von Neumann algebraic envelope of $A_r(w_r, t_r)$ will be isomorphic to the von Neumann algebraic envelope of $S_{q_r, \tau}^2$, which

is equal to $M_n(\mathbb{C})$, $B(l^2(\mathbb{N}))$ or $B(l^2(\mathbb{N})) \oplus B(l^2(\mathbb{N}))$, depending on whether $\varepsilon_r w_r$ is positive, zero or negative (cf. [28]). In any case, the corresponding image of Z_r will not contain 0 in its point-spectrum. ■

4 More on the $\varepsilon \in \text{Char}_{\{0,1\}}(Q^+)$ -case

The case of the non-standard Podleś spheres already shows that $W_\lambda(\mathfrak{g}; \varepsilon, +)$ is in general not a factor. However, for $\varepsilon \in \text{Char}_{\{0,1\}}(Q^+)$ and $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$ such that $W_\lambda(\mathfrak{g}; \varepsilon, +)$ is well-defined, we show that $W_\lambda(\mathfrak{g}; \varepsilon, +)$ *does* become a type I -factor, and we can then also say something more about the invariant integral $\varphi_\lambda^{\varepsilon,+}$ on $W_\lambda(\mathfrak{g}; \varepsilon, +)$.

Proposition 4.1. *Let $\varepsilon \in \text{Char}_{\{0,1\}}(Q^+)$, $\lambda \in \text{Char}_{\mathbb{R}^+}(P)$, and suppose $V_\lambda^{\varepsilon,+}$ is unitarizable with completion $\mathcal{H}_\lambda^{\varepsilon,+}$. Identify $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +) \subseteq B(\mathcal{H}_\lambda^{\varepsilon,+})$. Then this inclusion completes to a natural identification $W_\lambda(\mathfrak{g}; \varepsilon, +) \cong B(\mathcal{H}_\lambda^{\varepsilon,+})$.*

Proof. From the proof of Theorem 3.20 and Lemma 3.22, and from the commutation relations in Lemma 3.7, we get that the $A_r(w_r, t_r)$ appearing in the proof of Lemma 3.22 can only be matrix algebras or standard Podleś spheres. As the Z_r in the von Neumann algebraic completion of these algebras are always positive operators, it follows by induction that the components appearing in $W_\lambda(\mathfrak{g}; \varepsilon, +)$ arise from restrictions of highest weight modules $V_{\lambda'}^{\varepsilon,+}$ of $U_q(\mathfrak{g}; \varepsilon, +)$ with $\lambda' \in \text{Char}_{\mathbb{R}^+}(P)$. Our aim is to show that necessarily $\lambda' = \lambda$.

Let S be the set of r with $\varepsilon_r = 1$. Suppose that $\lambda' \in \text{Char}_{\mathbb{R}^+}(P)$ is such that the representation of $U_q(\mathfrak{g}; \varepsilon, +)$ on $V_{\lambda'}^{\varepsilon,+}$ factors over $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$. Suppose that $V_{\lambda'}^{\varepsilon,+}$ then admits an invariant pre-Hilbert space structure as a $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ -module, hence as a $U_q(\mathfrak{g}; \varepsilon, +)$ -module by Proposition 3.2. From the proof of Proposition 3.12, we deduce that there exists $w \in W_S$ such that $q^{(-2w\omega, \rho)} \lambda_{-4w\omega} = q^{(-2\omega, \rho)} \lambda'_{-4\omega}$ for all $\omega \in P$. In particular, $\lambda_{\omega_r} = \lambda'_{\omega_r}$ for $r \notin S$.

On the other hand, let $\tilde{U}_q(\mathfrak{k}_{\text{ss}})$ be the subalgebra of $U_q(\mathfrak{g}; \varepsilon, +)$ generated by the X_r^\pm and $K_{\alpha_r}^{\pm 1}$ with $r \in S$. Let $\tilde{V}_\lambda^{\varepsilon,+}$ be the $\tilde{U}_q(\mathfrak{k}_{\text{ss}})$ -module spanned by $v_\lambda^{\varepsilon,+}$, and similarly for $V_{\lambda'}^{\varepsilon,+}$. Then by Proposition 2.6, these are irreducible highest weight modules for $U_q(\mathfrak{k}_{\text{ss}})$ associated to the restrictions of λ and λ' to the root lattice Q_S of \mathfrak{k}_{ss} . But as $V_{\lambda'}^{\varepsilon,+}$ admits an invariant pre-Hilbert space structure (and \mathfrak{k}_{ss} is compact), it is necessarily finite-dimensional. However, as the restriction of λ' lies in the W_S -orbit of λ (for the so-called ‘dot’-action), it is well-known that this can happen only if the restrictions of λ and λ' to Q_S coincide. Combined with the observation at the end of the previous paragraph, this forces $\lambda = \lambda'$ inside $\text{Char}_{\mathbb{R}^+}(P)$. ■

Proposition 4.2. *Under the assumptions of Proposition 4.1, the invariant state on $W_\lambda(\mathfrak{g}; \varepsilon, +)$ is given by*

$$\varphi_\lambda^{\varepsilon,+}(x) = \frac{\text{Tr}(xZ_\rho)}{\text{Tr}(Z_\rho)},$$

where $Z_\rho = \prod_{r=1}^l Z_r$.

Proof. Consider the projection $E^{\varepsilon,+}$ of $U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$ onto its direct summand $\mathcal{L}(U_q(\mathfrak{g}; \varepsilon, +))$, the space of \triangleleft -invariants. For $\varepsilon_r > 0$ for all r , it follows from [16, Chapter 7] that $E^{\varepsilon,+}(xy) = E^{\varepsilon,+}(y\sigma(x))$ for all $x, y \in U_q(\mathfrak{g}; \varepsilon, +)_{\text{fin}}$, where $\sigma(x) = K_{-4\rho} x K_{-4\rho}^{-1}$. By continuity, this then holds for all $\varepsilon \in \text{Char}_{\mathbb{R}^+}(Q^+)$. Clearly σ induces an automorphism of $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$, which we will denote by the same symbol. As $\varphi_\lambda^{\varepsilon,+}$ factors over $E^{\varepsilon,+}$, we have as well that $\varphi_\lambda^{\varepsilon,+}(xy) = \varphi_\lambda^\varepsilon(y\sigma(x))$.

However, by general theory [3] we know that the modular automorphism group σ_t of $\varphi_\lambda^{\varepsilon,+}$ leaves $B_\lambda^{\text{fin}}(\mathfrak{g}; \varepsilon, +)$ invariant and is diagonalizable (and hence analytic) on it. From the above,

we can hence conclude that $\sigma_t = \text{Ad}(Z_\rho^{it})$ (where we use that Z_ρ is a positive operator by Proposition 4.1). As $W_\lambda(\mathfrak{g}; \varepsilon, +)$ is a type I -factor, again by Proposition 4.1, we know that $\varphi_\lambda^{\varepsilon, +}$ is completely determined by its modular automorphism group up to a scalar. Hence we obtain our expression for $\varphi_\lambda^{\varepsilon, +}$ as in the statement of the proposition. \blacksquare

A Cogroupoids

In this section, we recall the notion of cogroupoid due to J. Bichon [1, 2] (cf. also the notion of *face algebra* [13]).

Definition A.1. Let I be an index set, and let $\{H_{ij} \mid i, j \in I\}$ be a collection of $*$ -algebras. Suppose that for each triple of indices $i, j, k \in I$, we are given a unital $*$ -homomorphism

$$\Delta_{ij}^k: H_{ij} \rightarrow H_{ik} \otimes H_{kj}, \quad h \mapsto h_{(1)ik} \otimes h_{(2)kj}.$$

We call $(\{H_{ij}\}, \{\Delta_{ij}^k\})$ a connected *cogroupoid* over the index set I if the following conditions are satisfied:

- (Connectedness) None of the H_{ij} are the zero algebra.
- (Coassociativity) For each quadruple i, j, k, l of indices, we have

$$(\text{id} \otimes \Delta_{jl}^k) \Delta_{il}^j = (\Delta_{ik}^j \otimes \text{id}) \Delta_{il}^k,$$

- (Counits) There exist unital $*$ -homomorphisms $\epsilon_i: H_{ii} \rightarrow \mathbb{C}$ such that, for all indices i, j ,

$$(\epsilon_i \otimes \text{id}) \Delta_{ij}^i = \text{id}_{H_{ij}} = (\text{id} \otimes \epsilon_j) \Delta_{ij}^j,$$

- (Antipodes) There exist anti-homomorphisms $S_{ij}: H_{ij} \rightarrow H_{ji}$ such that, for all indices i, j and all $h \in H_{ii}$, we have

$$S_{ij}(h_{(1)ij})h_{(2)ji} = \epsilon_i(h) = h_{(1)ij}S_{ji}(h_{(2)ji}).$$

As for Hopf $*$ -algebras, it is easy to show that the S_{ij} are unique, and that $S_{ji}(S_{ij}(h)^*)^* = h$ for each $h \in H_{ij}$. Note that each (H_{ii}, Δ_{ii}^i) defines a Hopf $*$ -algebra.

Definition A.2. Let (H_{ij}, Δ_{ij}^k) be a cogroupoid. The *right adjoint action* (or Miyashita–Ulbrich action) \triangleleft of H_{jj} on H_{ij} is given by

$$x \triangleleft h = S_{ji}(h_{(1)ji})xh_{(2)ij}.$$

One easily proves that \triangleleft defines a right H_{jj} -module $*$ -algebra structure on H_{ij} . The compatibility with the $*$ -structure means that $(x \triangleleft h)^* = x^* \triangleleft S_{jj}(h)^*$.

Lemma A.3. *The space of \triangleleft -invariant elements in H_{ij} coincides with the center of H_{ij} .*

Proof. (Cf. [16, Lemma 2.4].) If $x \in H_{ij}$ is in the center, clearly $x \triangleleft h = \varepsilon_j(h)x$ for all $h \in H_{jj}$, by definition of the antipode. Conversely, if $S_{ji}(h_{(1)ji})xh_{(2)ij} = \varepsilon_j(h)x$ for all $h \in H_{jj}$, we have for $y \in H_{ij}$ that $xy = y_{(1)ij}S_{ji}(y_{(2)ji})xy_{(3)ij} = yx$. \blacksquare

Suppose now that B is a unital $*$ -algebra, and that for some i, j we have a unital $*$ -homomorphism $\pi: H_{ij} \rightarrow B$. As the right action of H_{jj} on H_{ij} is inner, it descends to a right action on B ,

$$b \triangleleft h = \pi(S_{ji}(h_{(1)ji}))b\pi(h_{(2)ij}).$$

Note that the central elements in B are invariant for the action.

A particular case of cogroupoid can be constructed from a Hopf algebra (H, Δ) together with a collection of real 2-cocycle functionals $\{\omega_i \mid i \in I\}$ on it. Here we mean by real 2-cocycle functional an element $\omega \in (H \otimes H)^*$ which is convolution invertible and such that $\omega(1, h) = \varepsilon(h) = \omega(h, 1)$ for all $h \in H$, while $\omega(h^*, k^*) = \overline{\omega(k, h)}$ and

$$\omega(h_{(1)}, k_{(1)})\omega(h_{(2)}k_{(2)}, l) = \omega(k_{(1)}, l_{(1)})\omega(h, k_{(2)}l_{(2)}), \quad \forall h, k, l \in H.$$

Let us write H_{ij} for the vector space H with the new multiplication

$$m_{ij}: H \otimes H \rightarrow H, \quad h \otimes k \mapsto \omega_i(h_{(1)}, k_{(1)})h_{(2)}k_{(2)}\omega_j^{-1}(h_{(3)}, k_{(3)}),$$

and write Δ_{ij}^k for the given coproduct Δ seen as a map $H_{ij} \rightarrow H_{ik} \otimes H_{kj}$. Then the (H_{ij}, Δ_{ij}^k) form a connected cogroupoid.

B Continuous one-parameter families of Lie algebras

We introduce the real Lie algebras whose quantizations we studied in Section 1. We refer to standard works as [14, 15, 22] for the basic background on Lie algebras and Lie groups.

We keep the notation as in Section 1. We further write $\{h_r\} \subseteq \mathfrak{h}$ for the basis dual to $\{\omega_r\}$, and write $\mathfrak{h}_{\mathbb{R}}$ for its real span, which we may identify with the real dual of $\mathfrak{h}_{\mathbb{R}}^*$. We write $\mathfrak{g}_c \subseteq \mathfrak{g}$ for the compact real form of \mathfrak{g} , and \dagger for the corresponding anti-linear anti-automorphism such that $x^\dagger = -x$ for $x \in \mathfrak{g}_c$.

Finally, for each $\alpha \in \Delta^+$, we choose root vectors $X_\alpha^\pm \in \mathfrak{n}^\pm$ such that $(X_\alpha^+)^\dagger = X_\alpha^-$ for all $\alpha \in \Delta^+$ and $[X_r^+, X_r^-] = h_r$ for all $\alpha_r \in \Phi^+$.

Fix $\varepsilon \in \text{Char}_{\mathbb{R}}(Q^+)$. We can define on \mathfrak{g} the linear maps

$$S_\varepsilon^\pm: \mathfrak{g} \rightarrow \mathfrak{g}: \begin{cases} X_\alpha^\pm \mapsto \varepsilon_\alpha X_\alpha^\pm, & \alpha \in \Delta^+, \\ z \mapsto z, & z \in \mathfrak{h} + \mathfrak{n}^\mp. \end{cases}$$

Definition B.1. We define \mathfrak{g}_ε to be the complex vector space

$$\{(S_\varepsilon^+(z), S_\varepsilon^-(z)) \mid z \in \mathfrak{g}\} \subseteq \mathfrak{g} \oplus \mathfrak{g}.$$

Proposition B.2. *The vector space \mathfrak{g}_ε is a Lie $*$ -subalgebra of the direct sum Lie algebra $\mathfrak{g} \oplus \mathfrak{g}$ equipped with the involution $(w, z)^* = (z^\dagger, w^\dagger)$. Moreover, $\dim_{\mathbb{C}}(\mathfrak{g}_\varepsilon) = \dim_{\mathbb{C}}(\mathfrak{g})$.*

Proof. From the definition, we see that \mathfrak{g}_ε is the linear space generated by elements of the form $(\varepsilon_\alpha X_\alpha^+, X_\alpha^+)$, $(X_\alpha^-, \varepsilon_\alpha X_\alpha^-)$ and (h_r, h_r) . Accordingly, \mathfrak{g}_ε has dimension $\dim_{\mathbb{C}}(\mathfrak{g})$ and inherits the $*$ -operation from $\mathfrak{g} \oplus \mathfrak{g}$. Since $\varepsilon_{\alpha+\beta} = \varepsilon_\alpha \varepsilon_\beta$ whenever α, β and $\alpha + \beta$ are positive roots, we find that \mathfrak{g}_ε is closed under the bracket operation. \blacksquare

In the following, we consider \mathfrak{g}_ε with its $*$ -operation inherited from $\mathfrak{g} \oplus \mathfrak{g}$.

Remark B.3. By rescaling the X_α^\pm , we see that $\mathfrak{g}_\varepsilon \cong \mathfrak{g}_\eta$ as Lie $*$ -algebras whenever $\varepsilon_r = \lambda_r \eta_r$ for certain $\lambda_r > 0$. Hence we may in principle always assume that $\varepsilon \in \{-1, 0, 1\}^l$. It is however sometimes convenient to keep the continuous deformation aspect into the game. In the physics literature, this type of deformation goes by the name of ‘the contraction method’. (In low dimensions, it can easily be visualized, cf. [12, Chapter 13].)

Definition B.4. We define $\mathfrak{g}_\varepsilon^{\mathbb{R}} = \{z \in \mathfrak{g}_\varepsilon \mid z^* = -z\}$.

Hence $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ is a real Lie algebra with \mathfrak{g}_ε as its complexification. We can also realize $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ more conveniently inside \mathfrak{g} as follows.

Proposition B.5. *Write*

$$X_\alpha^{(\varepsilon)} = X_\alpha^+ - \varepsilon_\alpha X_\alpha^-, \quad Y_\alpha^{(\varepsilon)} = i(X_\alpha^+ + \varepsilon_\alpha X_\alpha^-)$$

as elements in \mathfrak{g} . Consider the \mathbb{R} -linear span of the $X_\alpha^{(\varepsilon)}$, $Y_\alpha^{(\varepsilon)}$ and ih_r . Then this space is closed under the Lie bracket of \mathfrak{g} , and forms a real Lie algebra isomorphic to $\mathfrak{g}_\varepsilon^{\mathbb{R}}$.

Proof. As a real Lie algebra, one can embed \mathfrak{g} inside the direct sum $\mathfrak{g} \oplus \mathfrak{g}$ by means of the map

$$\mathfrak{g} \rightarrow \mathfrak{g} \oplus \mathfrak{g}, \quad z \mapsto (z, -z^\dagger).$$

Under this identification, it is immediately verified that the elements in the statement of the proposition get sent to a basis of $\mathfrak{g}_\varepsilon^{\mathbb{R}}$. \blacksquare

Proposition B.6. *Suppose $\varepsilon_r \neq 0$ for all r . Then $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ is a semi-simple real Lie algebra.*

Proof. In this case, the projection onto the first coordinate of $\mathfrak{g} \oplus \mathfrak{g}$ provides an isomorphism between \mathfrak{g}_ε and \mathfrak{g} . \blacksquare

Hence if $\varepsilon \in \text{Char}_{\{-1,1\}}(Q^+)$, the Lie algebra $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ is the real form of \mathfrak{g} corresponding to the involution

$$(X_r^+)^* = \varepsilon_r X_r^-, \quad h_r^* = h_r.$$

Accordingly, $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ is a real Lie algebra of equal rank, meaning that we can take a Cartan decomposition $\mathfrak{g}^{\mathbb{R}} = \mathfrak{t} \oplus \mathfrak{p}$ with \mathfrak{t} compact and $i\mathfrak{h}_{\mathbb{R}} \subseteq \mathfrak{t}$. Conversely, it is easy to see that any equal rank semi-simple real Lie algebra can be realized as some $\mathfrak{g}_\varepsilon^{\mathbb{R}}$. We recall that the equal rank semi-simple real Lie algebras are precisely those which admit an irreducible unitary representation of discrete type.

As an example, let us present the simple equal rank real Lie algebras of type (A) (see [14, Chapter X]). For $r \in \{0, \dots, l\}$, choose $\sigma_r \in \{\pm 1\}$ such that $\varepsilon_r = \sigma_{r-1}\sigma_r$ for $r \in \{1, \dots, l\}$. Then $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ is always of type (AIII), and ε corresponds to $\mathfrak{su}(p, l+1-p)$ with p the number of negative σ_r . This precise correspondence between the ε and the various real Lie algebras is easy to determine explicitly from the standard descriptions of the Cartan decompositions.

More generally, we obtain from Proposition B.5 the following characterization of which Lie algebras appear as some \mathfrak{g}_ε .

Corollary B.7. *Let $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$ be a parabolic Lie subalgebra of \mathfrak{g} with Levi factor \mathfrak{l} and largest nilpotent ideal \mathfrak{u} . Let $\mathfrak{l}^{\mathbb{R}}$ be a real form of \mathfrak{l} which is of equal rank on each simple summand and compact on the center. Then $\mathfrak{l}^{\mathbb{R}} \oplus \mathfrak{u} \cong \mathfrak{g}_\varepsilon^{\mathbb{R}}$ as real Lie algebras for some $\varepsilon \in \text{Char}_{\mathbb{R}}(Q^+)$, and all $\mathfrak{g}_\varepsilon^{\mathbb{R}}$ arise in this way.*

Our next step is to present \mathfrak{g}_ε by means of generators and relations.

Definition B.8. Fix $\varepsilon \in \text{Char}_{\mathbb{R}}(Q^+)$. We define $\tilde{\mathfrak{g}}_\varepsilon$ as a universal Lie algebra in the following way. A set of generators is given by a triple of elements X_r^\pm and H_r for each simple root α_r . The relations can be grouped into four parts, which we label as ‘(H)-condition’, ‘(T)orus action’, ‘(S)erre relations’ and ‘(C)oupling conditions’: for all $r, s \in I$,

$$(H) \quad [H_r, H_s] = 0,$$

$$(T) \quad [H_r, X_s^\pm] = \pm a_{rs} X_s^\pm,$$

$$(S) \quad \text{ad}(X_r^\pm)^{1-a_{rs}}(X_s^\pm) = 0 \text{ when } r \neq s,$$

$$(C)_\varepsilon \quad [X_r^+, X_s^-] = \delta_{rs} \varepsilon_r H_r.$$

It is immediate that $\tilde{\mathfrak{g}}_\varepsilon$ can be endowed with a $*$ -operation such that $(X_r^+)^* = X_r^-$ and $H_r^* = H_r$.

Proposition B.9. *The Lie $*$ -algebras $\tilde{\mathfrak{g}}_\varepsilon$ and \mathfrak{g}_ε are isomorphic.*

Proof. It is straightforward to verify that there is a unique Lie $*$ -algebra homomorphism $\phi: \tilde{\mathfrak{g}}_\varepsilon \rightarrow \mathfrak{g}_\varepsilon \subseteq \mathfrak{g} \oplus \mathfrak{g}$ such that $\phi(X_r^+) = (\varepsilon_r X_r^+, X_r^+)$, $\phi(X_r^-) = (X_r^-, \varepsilon_r X_r^-)$ and $\phi(H_r) = (h_r, h_r)$. It is obviously surjective. On the other hand, it is easy to see by induction that each element of $\tilde{\mathfrak{g}}_\varepsilon$ can be written in the form $x + y + z$ with x in the Lie algebra generated by the X_r^+ 's, z in the Lie algebra generated by the X_r^- 's, and y in the linear span of the H_r . Since \mathfrak{n}^\pm are universal with respect to the relations (S), we find that $\dim(\tilde{\mathfrak{g}}_\varepsilon) \leq \dim(\mathfrak{g}_\varepsilon)$, hence ϕ is bijective. ■

Acknowledgments

It is a pleasure to thank the following people for discussions on topics related to the subject of this paper: J. Bichon, P. Bieliavsky, H.P. Jakobsen, E. Koelink, S. Kolb, U. Krähmer and S. Neshveyev.

References

- [1] Bichon J., Hopf–Galois systems, *J. Algebra* **264** (2003), 565–581, [math.QA/0204348](#).
- [2] Bichon J., Hopf–Galois objects and cogroupoids, [arXiv:1006.3014](#).
- [3] Boca F.P., Ergodic actions of compact matrix pseudogroups on C^* -algebras, *Astérisque* **232** (1995), 93–109.
- [4] Chari V., Pressley A., A guide to quantum groups, Cambridge University Press, Cambridge, 1995.
- [5] De Commer K., Comonoidal W^* -Morita equivalence for von Neumann bialgebras, *J. Noncommut. Geom.* **5** (2011), 547–571, [arXiv:1004.0824](#).
- [6] De Commer K., On a correspondence between $SU_q(2)$, $\tilde{E}_q(2)$ and $\widetilde{S}U_q(1, 1)$, *Comm. Math. Phys.* **304** (2011), 187–228, [arXiv:1004.4307](#).
- [7] De Commer K., On a Morita equivalence between the duals of quantum $SU(2)$ and quantum $\tilde{E}(2)$, *Adv. Math.* **229** (2012), 1047–1079, [arXiv:0912.4350](#).
- [8] De Commer K., On the construction of quantum homogeneous spaces from $*$ -Galois objects, *Algebr. Represent. Theory* **15** (2012), 795–815, [arXiv:1001.2153](#).
- [9] De Concini C., Kac V.G., Representations of quantum groups at roots of 1, in Operator Algebras, Unitary Representations, Enveloping Algebras, and Invariant Theory (Paris, 1989), *Progr. Math.*, Vol. 92, Birkhäuser Boston, Boston, MA, 1990, 471–506.
- [10] Donin J., Mudrov A., Explicit equivariant quantization on coadjoint orbits of $GL(n, \mathbb{C})$, *Lett. Math. Phys.* **62** (2002), 17–32, [math.QA/0206049](#).
- [11] Enock M., Morita equivalence of measured quantum groupoids. Application to deformation of measured quantum groupoids by 2-cocycles, in Operator Algebras and Quantum Groups, *Banach Center Publ.*, Vol. 98, Editors W. Pusz, P.M. Soltan, Polish Acad. Sci. Inst. Math., Warsaw, 2012, 107–198, [arXiv:1106.1018](#).
- [12] Gilmore R., Lie groups, physics, and geometry. An introduction for physicists, engineers and chemists, [Cambridge University Press](#), Cambridge, 2008.
- [13] Hayashi T., Face algebras. I. A generalization of quantum group theory, *J. Math. Soc. Japan* **50** (1998), 293–315.
- [14] Helgason S., Differential geometry, Lie groups, and symmetric spaces, *Pure and Applied Mathematics*, Vol. 80, Academic Press Inc., New York, 1978.
- [15] Humphreys J.E., Representations of semisimple Lie algebras in the BGG category \mathcal{O} , *Graduate Studies in Mathematics*, Vol. 94, American Mathematical Society, Providence, RI, 2008.
- [16] Joseph A., Letzter G., Local finiteness of the adjoint action for quantized enveloping algebras, *J. Algebra* **153** (1992), 289–318.
- [17] Joseph A., Letzter G., Separation of variables for quantized enveloping algebras, *Amer. J. Math.* **116** (1994), 127–177.

- [18] Joseph A., Todoric D., On the quantum KPRV determinants for semisimple and affine Lie algebras, *Algebr. Represent. Theory* **5** (2002), 57–99.
- [19] Karolinsky E., Stolin A., Tarasov V., Irreducible highest weight modules and equivariant quantization, *Adv. Math.* **211** (2007), 266–283, [math.QA/0507348](#).
- [20] Kashiwara M., On crystal bases of the Q -analogue of universal enveloping algebras, *Duke Math. J.* **63** (1991), 465–516.
- [21] Kassel C., Schneider H.J., Homotopy theory of Hopf Galois extensions, *Ann. Inst. Fourier (Grenoble)* **55** (2005), 2521–2550, [math.QA/0402034](#).
- [22] Knapp A.W., Representation theory of semisimple groups. An overview based on examples, *Princeton Landmarks in Mathematics*, Princeton University Press, Princeton, NJ, 2001.
- [23] Knapp A.W., Lie groups beyond an introduction, 2nd ed., *Progress in Mathematics*, Vol. 140, [Birkhäuser Boston, Inc.](#), Boston, MA, 2002.
- [24] Korogodsky L.I., Representations of quantum algebras arising from non-compact quantum groups: Quantum orbit method and super-tensor products, Ph.D. Thesis, Massachusetts Institute of Technology, 1996, available at <http://dspace.mit.edu/handle/1721.1/39076>.
- [25] Letzter G., Quantum symmetric pairs and their zonal spherical functions, *Transform. Groups* **8** (2003), 261–292, [math.QA/0204103](#).
- [26] Letzter G., Quantum zonal spherical functions and Macdonald polynomials, *Adv. Math.* **189** (2004), 88–147, [math.QA/0210447](#).
- [27] Levendorskiĭ S., Soibelman Y., Algebras of functions on compact quantum groups, Schubert cells and quantum tori, *Comm. Math. Phys.* **139** (1991), 141–170.
- [28] Masuda T., Nakagami Y., Watanabe J., Noncommutative differential geometry on the quantum two sphere of Podleś. I. An algebraic viewpoint, *K-Theory* **5** (1991), 151–175.
- [29] Mudrov A., Quantum conjugacy classes of simple matrix groups, *Comm. Math. Phys.* **272** (2007), 635–660, [math.QA/0412538](#).
- [30] Reshetikhin N., Yakimov M., Quantum invariant measures, *Comm. Math. Phys.* **224** (2001), 399–426, [math.QA/0101048](#).
- [31] Rosso M., Groupes quantiques, représentations linéaires et applications, Ph.D. Thesis, Université de Paris VII, 1990.
- [32] Vaksman L.L., Quantum bounded symmetric domains, *Translations of Mathematical Monographs*, Vol. 238, American Mathematical Society, Providence, RI, 2010.
- [33] Woronowicz S.L., Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665.
- [34] Woronowicz S.L., Twisted $SU(2)$ group. An example of a noncommutative differential calculus, *Publ. Res. Inst. Math. Sci.* **23** (1987), 117–181.